



Fluro IndoBlu cell proliferation Detection Kit

Product Information

Cat.No.

Kit-0204

Description

Measurement of cell proliferation and viability is frequently used in clinical and experimental immunology as means of assessing cell activation in response to diseases, infections and environmental stimulations. Fluoro IndoBlu assay can be used for the measurement of cell proliferation in response to antigens, cytokines, growth factors, and mitogens. It can also be used for the analysis of cytotoxic effects of anticancer drugs, drug resistance, cytotoxic pharmaceutical compounds, and other toxic agents. Staurosporine induced Jurkat cell viability assay. Titrated doses of staurosporine were added into seeded Jurkat (10,000 cells/well) cells in 96 well black opaque tissue culture plates and incubated at 37°C, 10% CO₂ for 24h. Fluoro IndoBlu™ (1/10 volume) was added into the cultured cells and incubated for 3h and fluorescence was detected at Ex: 530nm and Em: 590nm. PBMC proliferation in response to mitogen Concanavalin A (Con A) stimulation measured by Fluoro IndoBlu™. PBMC at 15,000 cells were cultured for 3 days in the presence of titrated amounts of Con-A and tested for proliferation using the Fluoro IndoBlu™ assay. Fluoro IndoBlu™ (1/10 volume) was added to the cultured cells and incubated 37°C for 2h. Fluorescence was detected at Ex: 530nm and Em: 590nm.

Applications

Fluorescence plate reader

Usage

1. This kit is only for usage on cell lines or primary cell cultures. 2. For research use only. Not for use in diagnostic procedures. 3. Gloves, protective clothing and eyewear should be worn and safe laboratory practices followed.

Storage

1. Short term: Store the kit at 2-8°C for (1 -2 weeks), keep it away from light. 2. Long term: Store the kit at -20°C.



CREATIVE BIOMART[®]
Assay Kit

Fluro IndoBlu cell proliferation Detection Kit

Kit Components

1. IndoBlu Reagent: 2-8°C; 2. Optional: 96-Well Clear Bottomed Black Plate for Fluorescent Readout

Features & Benefits

1. Highly effective and stable assay for cell proliferation. 2. No need to wash out media from cell samples, just add the reagent directly to your experimental samples. 3. Works for both cell viability and proliferation assay. 4. Plate can be read in 30 minutes to 3 hours.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.org

Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA