

DNAse I Activity Assay Kit (Fluorometric)

Product Information

Cat

Kit-1036

Cat.No.

Kit-1036

Description

Deoxyribonuclease I (DNAse I) is an endonuclease that cleaves DNA phosphodiester bonds yielding 5'-phosphorylated and 3'-hydroxylated oligonucleotides. DNAse I targets single-stranded DNA, double-stranded DNA, and chromatin in a non-specific manner. As an important player in cellular waste management, DNAse I is normally secreted extracellularly to clear the system from circulating cell-free DNA, foreign DNA from food digestion or potential pathogens, and endogenous chromosomal DNA from apoptotic and necrotic cells. Abnormal DNAse I activity occurs in association with a variety of cancers and auto-immune illnesses that exhibit elevated levels of cell-free DNA. Furthermore, DNAse I has been therapeutically used in cystic fibrosis patients to degrade DNA and reduce sputum viscosity. DNAse I Activity Fluorometric Assay Kit allows for quantitative evaluation of DNAse I activity of purified enzymes and their inhibitors as well as comparative examination of DNAse I activity in biological samples. Enzyme activity is detected upon cleavage of a DNA Probe, which yields a fluorescent DNA product measured at Ex/Em = 651/681 nm. The limit of quantification (L.O.Q) is 178 fmoles of DNA probe cleaved per minute per ml.

Applications

Measurement of DNAse I activity of purified proteins;

Quantitative analysis of DNAse I mutants and inhibitors;

Comparative examination of DNAse I activity in biological samples

Storage

Store kit at -20°C, protected from light. Briefly centrifuge small vials prior to opening. Upon re-suspension, aliquot DNA Probe and DNAse I Positive control to avoid repeated freeze-thaw cycles. Read entire protocol before performing the assay.

DNAse I Activity Assay Kit (Fluorometric)

- 10X DNAse I Assay Buffer: Store at 4 °C. Warm to 37 °C temperature before use.
- DNA Probe: Reconstitute with 220 µl of DNA Probe Re-suspension Buffer. Aliquot and store at -20°C. Avoid multiple freeze-thaw cycles.
- DNA Probe Re-suspension Buffer: Ready to use. Store at RT.
- DNAse I Positive Control: Reconstitute with 220 µl of Positive Control Re-suspension Buffer. Aliquot and store at -20°C.
- Positive Control Re-suspension Buffer: Ready to use. Store at -20°C.
- Molecular Biology Grade Water: Ready to use. Store at RT.

Size

100 assays

Kit Components

10X DNAse I Assay Buffer, 1.1 ml

DNA Probe, 1 vial

DNA Probe Re-suspension Buffer, 250 µl

DNAse I Positive Control, 1 vial

Positive Control Re-suspension Buffer, 1 ml

Molecular Biology Grade Water, 25 ml

Materials Required but Not Supplied

96-well white plate with flat bottom, low-medium binding

Spectrophotometer

Purified DNAse I, DNAse I inhibitors, biological samples

Compatible Sample Types

Purified Protein, serum.

Assay Protocol

Caution! It is imperative to use molecular biology grade water for sample preparation and filter tips for sample pipetting at all times to avoid DNAse contamination.

1. Sample Preparation: Thaw any purified enzymes and biological samples along with all the provided assay components on ice, unless otherwise stated. Dilute enzymes, inhibitors, and

DNAse I Activity Assay Kit (Fluorometric)

biological samples to a desired concentration with water or their corresponding storage buffer. Add a desired amount of enzyme, inhibitor, or biological sample to each well and adjust the volume to 50 μ l with water. Use water only (no enzyme/sample) for background control reaction. For positive control reaction, add 2 μ l of DNAse I Positive Control to 48 μ l of water. Mix well.

Notes:

- a. Do not store enzyme/inhibitor/sample dilutions; discard the dilutions instead.
- b. The recommended amount of serum sample to use in the assay is 10-25 μ l.
- c. For uncharacterized enzymes, we suggest testing several doses to ensure the reading is within the Standard Curve range.
- d. If the user suspects any non-specific sample DNAse activity, 50 mM 2-Nitro-5-thiocyanatobenzoic acid can be used to specifically inhibit DNAse I activity.

2. DNA Probe Standard Curve: Prepare 1 μ M DNA Probe stock by diluting 4 μ l of 25 μ M DNA Probe in 96 μ l of molecular biology grade water. Add 0, 4, 8, 12, 16, 20 μ l of 1 μ M DNA Probe into a series of wells on a 96-well plate to generate 0, 4, 8, 12, 16, 20, pmol/well of DNA Probe Standard. Adjust the volume to 50 μ l with molecular biology grade water.

3. Reaction Mix: Mix enough reagents for the number of assays to be performed. For each well containing sample(s), and standards prepare 50 μ l Mix containing:

Sample Reaction Mix DNA Probe Standard Reaction Mix

(1 assay) (1 assay)

10X DNAse I Assay Buffer 10 μ l 10 μ l

DNA Probe (25 μ M) 2 μ l ---

DNAse I Positive Control --- 2 μ l

Molecular Biology Grade H₂O 38 μ l 38 μ l

Mix and add 50 μ l of the Sample Reaction Mix to each well containing the Positive Control, Test Samples, and Background Control. Add 50 μ l of DNA Probe Standard Reaction Mix to each well containing DNA Probe Standard.

4. Measurement: For positive control, test samples, background control, and DNA Probe Standard measure fluorescence (Ex/Em = 651/681 nm) in kinetic mode every 30 seconds for at least 90 minutes at 37 °C. Adjust GAIN/PMT setting of your fluorometer as necessary so that the standard

DNAse I Activity Assay Kit (Fluorometric)

curve readings are within the detection range of the instrument.

Analysis

5. Calculations: Standard Curve: Record RFU at $t = 90$ min for each DNA Probe standard curve reading. Plot the DNA Probe standard curve with pmol of DNA on the x-axis and RFU on the y-axis. Apply a linear fit to the DNA standard values and determine the standard curve equation. Samples/Positive Control: Apply RFU values at each time point to the standard curve equation to determine pmol of DNA cleaved at each reaction time point. Plot pmol DNA on the y-axis vs. time (in minutes) on the x-axis and determine the slope (pmol/min) of the linear portion of the reaction curve. Subtract background control readings from samples.

Sample DNAse I Activity = (slope/V) \times D (pmol/min/ml \equiv μ U/ml)

Sample Specific Activity = (slope/ μ g) \times D (pmol/min/ μ g \equiv μ U/ μ g)

Where: V = sample volume added into the reaction well (ml).

D = Dilution Factor

Slope = pmol/min (from the linear range of the activity curve)

Unit Definition: One unit of DNAse I is the amount of enzyme that cleaved 1.0 μ mol of DNA Probe per min. at 37°C.