



Pan-Methyl Histone H3-K79 Quantification Kit (Colorimetric)

Product Information

Cat.No.

Kit-0655

Product Overview

Pan-Methyl Histone H3-K79 Quantification Kit (Colorimetric) is use for measuring mono, di-and tri-methylation of histone H3-K79.

Description

Epigenetic activation or inactivation of genes plays a critical role in many important human diseases, especially in cancer. A major mechanism for epigenetic inactivation of the genes is methylation of CpG islands in genome DNA caused by DNA methyltransferases. Histone methyltransferases (HMTs) control or regulate DNA methylation through chromatin-dependent transcription repression or activation. HMTs transfer 1-3 methyl groups from S-adenosyl-L-methionine to the lysine and arginine residues of histone proteins. Dot1 is a histone methyltransferase that catalyzes methylation of histone H3 at lysine 79 (H3-K79) in mammalian cells. Intergenic H3-K79 mono-methylation is a mark of enhancer regions and may provide unique binding surfaces along the chromatin fiber to stabilize interactions between enhancers and their target promoters. H3-K79 diand tri-methylation is also a widespread histone modification and is associated with transcriptionally active genes. Increased H3-K79 methylation is found to be involved in some pathological processes such as leukemogenesis in human. The patterns of H3-K79 methylation can also be changed by inhibition or activation of HMTs. Thus quantitative detection of mono-, di- and tri-methyl histone H3-K79 would provide useful information for better understanding epigenetic regulation of gene activation and for developing HMT targeted drugs. The Pan-Methyl Histone H3-K79 Quantification Kit (Colorimetric) provides a tool for measuring mono, di- and tri-methylation of histone H3-K79.

Applications

For simultaneously measuring histone H3-K79 mono-, di-, and tri-methylation using a variety of mammalian cells (human, mouse, etc.) including fresh and frozen tissues, cultured adherent and suspension cells.



Pan-Methyl Histone H3-K79 Quantification Kit (Colorimetric)

Usage

For research use only (RUO)

Storage

Upon receipt, store C3 and Standard control at -20°C . Store all other components at 4°C away from light. The components of the kit should be stable for 6 months when stored properly. Note: Check if buffers C1 and C2 contain salt precipitates before using. If so, warm (at room temperature or 37°C) and shake the buffers until the salts are redissolved.

Kit Components

C1 (10X wash buffer) 20 ml C2 (antibody buffer) 12 ml C3 (detection antibody, 1 mg/ml)* 10 μl C4 (color developer) 10 μl C5 (stop solution) 6 ml Standard control (100 $\mu\text{g}/\text{ml}$)* 20 μl 8 well sample strips (with frame) 98 well standard control strips* 3* For maximum recovery of the products, centrifuge the original vial after thawing prior to opening the cap.

Features & Benefits

Quick and efficient procedure, which can be finished within 2.5 hours. Innovative colorimetric assay with no need for radioactivity, electrophoresis, and chromatography. Simultaneous quantification of mono-, di-, and tri-methylated H3-K79 with the detection limit as low as 2 ng/well and detection range from 20 ng-5 $\mu\text{g}/\text{well}$ of histone extracts. The control is conveniently included for quantification of the amount of mono-, di- and tri-methylated H3-K79. Strip microplate format makes the assay flexible: manual or high throughput. Simple, reliable, and consistent assay conditions.
