



Pan-Methyl Histone H3-K9 Quantification Kit (Fluorometric)

Product Information

Cat.No.

Kit-0658

Product Overview

Pan-Methyl Histone H3-K9 Quantification Kit (Fluorometric) is use for measuring mono-, di-and tri-methylation of histone H3-K9.

Description

Epigenetic activation or inactivation of genes plays a critical role in many important human diseases, especially in cancer. A major mechanism for epigenetic inactivation of the genes is methylation of CpG islands in genome DNA caused by DNA methyltransferases. Histone methyltransferases (HMTs) control or regulate DNA methylation through chromatin-dependent transcription repression or activation. HMTs transfer 1-3 methyl groups from S-adenosyl-L-methionine to the lysine and arginine residues of histone proteins. ESET, G9a, SUV39-h1, SUV39- h2, SETDB1, Dim-5 and Eu-HMTase are histone methyltransferases that catalyze methylation of histone H3 at lysine 9 (H3-K9). In mammalian cells, mono-methyl H3-K9 are enriched in certain euchromatic domains, which have been postulated to be transcriptionally silent. H3-K9 di- and tri-methylation mediates heterochromatin formation by forming a binding site for HP1 and also participates in silencing gene expression at euchromatic sites. Increased H3-K9 methylation is also found to be involved in some pathological processes such as cancer progression. The patterns of H3-K9 methylation can be also changed by inhibition or activation of HMTs. Thus quantitative detection of the patterns of histone H3-K9 methylation would provide useful information for better understanding epigenetic regulation of gene activation/silencing and for developing HMT-targeted drugs. The Pan-Methyl Histone H3-K9 Quantification Kit (Colorimetric) provides a tool for measuring mono-, di- and tri-methylation of histone H3-K9.

Applications

For simultaneously measuring histone H3-K9 mono-, di-, and tri-methylation using a variety of mammalian cells (human, mouse, etc.) including fresh and frozen tissues, cultured adherent and suspension cells.



Pan-Methyl Histone H3-K9 Quantification Kit (Fluorometric)

Usage

For research use only (RUO)

Storage

Upon receipt, store F3, F4 and Standard control at -20°C . Store all other components at 4°C away from light. The components of the kit should be stable for 6 months when stored properly. Note: Check if buffers F1 and F2 contain salt precipitates before using. If so, warm (at room temperature or 37°C) and shake the buffers until the salts are redissolved.

Kit Components

F1 (10X wash buffer) 20 ml F2 (antibody buffer) 12 ml F3 (detection antibody, 1 mg/ml)* 10 μl F4 (fluoro-developer)* 24 μl F5 (fluoro enhancer)* 24 μl F6 (fluoro-dilution) 8 ml Standard control (100 $\mu\text{g}/\text{ml}$)* 20 μl 8 well sample strips (with frame) 98 well standard control strips* 3* For maximum recovery of the products, centrifuge the original vial after thawing prior to opening the cap.

Features & Benefits

Quick and efficient procedure, which can be finished within 3 hours. Innovative fluorometric assay with no need for radioactivity, electrophoresis, and chromatograph. Simultaneous quantification of mono-, di-, and tri-methylated H3-K9 with the detection limit as low as 0.4 ng/well and detection range from 5 ng-2 $\mu\text{g}/\text{well}$ of histone extracts. The control is conveniently included for quantification of the amount of mono-, di- and tri-methylated H3-K9. Strip microplate format makes the assay flexible: manual or high throughput. Simple, reliable, and consistent assay conditions.
