

Cyclic Nucleotide Phosphodiesterase assay Kit

Product Information

Cat.No.

Kit-0667

Product Overview

Unique, non-radioactive, colorimetric assay. Used to screen inhibitors and modulators of cyclic nucleotide phosphodiesterase activity. HTS format permitting rapid assays of large numbers of samples.

Size

96 wells

Description

A BIOMOL GREEN Quantizyme Assay System. The Cyclic Nucleotide Phosphodiesterase (PDE) Assay Kit combines a special dual enzyme system with the BIOMOL GREEN Reagent for phosphate detection to create a unique, non-radioactive, colorimetric assay. This HTS-friendly, mix and read system may be used to screen inhibitors and modulators of cyclic nucleotide phosphodiesterase activity. 96-well microplate format permits rapid assays of large numbers of samples. The basis for the assay is the cleavage of cAMP or cGMP by a cyclic nucleotide phosphodiesterase. The 5"-nucleotide released is further cleaved into the nucleoside and phosphate by the enzyme 5"-nucleotidase. The phosphate released due to enzymatic cleavage is quantified using BIOMOL GREEN reagent in a modified Malachite Green assay. The kit includes Type I cyclic AMP phosphodiesterase (PDE) for validation purposes.

Applications

Colorimetric detection, HTS

Storage

-80°C

Kit Components

PDE Enzyme (from bovine brain) (5 x 4 U; lyophilized solid, 4 U per vial. 1 U = 1 nmol 3",5-cAMP to

Cyclic Nucleotide Phosphodiesterase assay Kit

5"-AMP per minute under the conditions of the linearity assay (30°C, pH 7.4, 200µM 3",5"-cAMP). Storage: -70°C, avoid freeze/thaw cycles. 5"-Nucleotidase (1ml; 5 kU/µl in 10mM TRIS-hydrochloric acid, pH 7.4, 125mM sodium chloride, 10% glycerol, 1mM magnesium chloride. One U will release 1 pmol phosphate per minute from 200µM 5"-AMP, 30°C in a reaction buffer of 10mM TRIS-hydrochloric acid, pH 7.4, 0.2mM magnesium chloride). Storage: -70°C 3",5"-cAMP Substrate (2ml; 1mM in assay buffer (10mM TRIS-hydrochloric acid, pH 7.4)). Storage: -20 or -70°C 3",5"-cGMP Substrate (2ml; 1mM in assay buffer (10mM TRIS-hydrochloric acid, pH 7.4)). Storage: -20 or -70°C CPDE Assay Buffer (40ml (2 x 20ml); 10mM TRIS-hydrochloric acid, pH 7.4) Storage: -70°C Biomol Green Reagent (20ml; liquid in screw-cap plastic bottle). Storage: 4°C, long-term at -70°C 5"-AMP Standard (1.0ml; 100µM in 10mM TRIS-hydrochloric acid, pH 7.4 buffer). Storage: -20 or -70°C 5"-GMP Standard (1.0ml; 100µM in 10mM TRIS-hydrochloric acid, pH 7.4 buffer). Storage: -20 or -70°C Inhibitor (IBMX) (200µl; 200µM in assay buffer (10mM TRIS-hydrochloric acid, pH 7.4)). Storage: -20 or -70°C Desalting column and resin (1 column and 1 g of resin). Storage: room temperature, after rehydration store at 4°C 1/2 volume microplate. Storage: Room temperature