



# Tri-Methyl Histone H3-K4 Quantification Kit (Colorimetric)

## Product Information

---

### Cat.No.

Kit-0822

---

### Product Overview

Tri-Methyl Histone H3-K4 Quantification Kit (Colorimetric) is use for measuring tri-methylation of histone H3-K4.

---

### Description

Epigenetic activation or inactivation of genes plays a critical role in many important human diseases, especially in cancer. A major mechanism for epigenetic inactivation of the genes is methylation of CpG islands in genome DNA caused by DNA methyltransferases. Histone methyltransferases (HMTs) control or regulate DNA methylation through chromatin-dependent transcription repression or activation. HMTs transfer 1-3 methyl groups from S-adenosyl-Lmethionine to the lysine and arginine residues of histone proteins. SET1, SET7/9, Ash1, ALL-1, MLL, ALR, Trx, and SMYD3 are histone methyltransferases that catalyze methylation of histone H3 at lysine 4 (H3-K4) in mammalian cells. H3- K4 tri-methylation has been viewed as a signature mark of highly transcribed genes, which is placed exclusively in the 5'- region downstream of the promoter. Increased H3-K4 tri-methylation is also found to be involved in some pathological processes such as cancer progression. The H3-K4 trimethylation can be also changed by inhibition or activation of HMTs. Thus, quantitative detection of tri-methyl histone H3-K4 would provide useful information for better understanding epigenetic regulation of gene activation, and for developing HMT-targeted drugs.

---

### Applications

Tri-Methyl Histone H3-K4 Quantification Kit (Colorimetric) is suitable for specifically measuring histone H3-K4 tri-methylation using a variety of mammalian cells (human, mouse, etc.) including fresh and frozen tissues, and cultured adherent and suspension cells.

---

### Usage

For research use only (RUO)

---

### Storage

---



## Tri-Methyl Histone H3-K4 Quantification Kit (Colorimetric)

Upon receipt, store C3 and Standard control at  $-20^{\circ}\text{C}$ . Store all other components at  $4^{\circ}\text{C}$  away from light. The components of the kit should be stable for 6 months when stored properly. Note: Check if buffers C1 and C2 contain salt precipitates before using. If so, warm (at room temperature or  $37^{\circ}\text{C}$ ) and shake the buffers until the salts are redissolved.

### Kit Components

C1 (10X wash buffer) 20 ml C2 (Antibody buffer) 12 ml C3 (Detection Antibody, 1 mg/ml)\* 10  $\mu\text{l}$  C4 (Color Developer) 10 ml C5 (Stop Solution) 6 ml Standard Control (100  $\mu\text{g}/\text{ml}$ )\* 20  $\mu\text{l}$  8-well sample strips (with frame) 9 strips 8-well standard control strips\* 3 strips\* For maximum recovery of the products, centrifuge the original vial after thawing prior to opening the cap.

### Features & Benefits

Quick and efficient procedure, which can be finished within 2 hours 30 minutes. Innovative colorimetric assay with no need for radioactivity, electrophoresis, and chromatography. Specifically capturing tri-methylated H3-K4 with the detection limit as low as 2 ng/well and detection range from 20 ng-5  $\mu\text{g}/\text{well}$  of histone extracts. The control is conveniently included for quantification of the amount of tri-methylated H3-K4. Strip microplate format makes the assay flexible: manual or high throughput. Simple, reliable, and consistent assay conditions.