



Tri-Methyl Histone H3-K79 Quantification Kit (Fluorometric)

Product Information

Cat.No.

Kit-0825

Product Overview

Tri-Methyl Histone H3-K79 Quantification Kit (Fluorometric) is use for measuring tri-methylation of histone H3-K79.

Description

Epigenetic activation or inactivation of genes plays a critical role in many important human diseases, especially in cancer. A major mechanism for epigenetic inactivation of the genes is methylation of CpG islands in genome DNA caused by DNA methyltransferases. Histone methyltransferases (HMTs) control or regulate DNA methylation through chromatin-dependent transcription repression or activation. HMTs transfer 1-3 methyl groups from S-adenosyl-L-methionine to the lysine and arginine residues of histone proteins. Dot1 is a histone methyltransferase that catalyzes methylation of histone H3 at lysine 79 (H3-K79) in mammalian cells. H3-K79 tri-methylation is widespread histone modification and associated with transcriptionally active genes. Increased H3-K79 trimethylation is also found to be involved in some pathological processes such as leukemogenesis in human. The H3-K79 tri-methylation can be also changed by inhibition or activation of HMTs. Thus quantitative detection of tri-methyl histone H3-K79 would provide useful information for better understanding epigenetic regulation of gene activation and for developing HMT targeted drugs. The Tri-Methyl Histone H3-K79 Quantification Kit (Colorimetric) provides a tool for measuring tr-methylation of histone H3- K79.

Applications

For specifically measuring histone H3-K79 trimethylation using a variety of mammalian cells (human, mouse, etc.) including fresh and frozen tissues, cultured adherent and suspension cells.

Usage

For research use only (RUO)

Storage



Tri-Methyl Histone H3-K79 Quantification Kit (Fluorometric)

Upon receipt, store F3, F4 and Standard control at -20°C . Store all other components at 4°C away from light. The components of the kit should be stable for 6 months when stored properly. Note: Check if buffers F1 and F2 contain salt precipitates before using. If so, warm (at room temperature or 37°C) and shake the buffers until the salts are redissolved.

Kit Components

F1 (10X wash buffer) 20 ml F2 (antibody buffer) 12 ml F3 (detection antibody, 1 mg/ml)* 10 μl F4 (fluoro-developer)* 24 μl F5 (fluoro enhancer)* 24 μl F6 (fluoro-dilution) 8 ml Standard control (100 $\mu\text{g}/\text{ml}$)* 20 μl 8 well sample strips (with frame) 98 well standard control strips* 3* For maximum recovery of the products, centrifuge the original vial after thawing prior to opening the cap.

Features & Benefits

Quick and efficient procedure, which can be finished within 2.5 hours. Innovative fluorometric assay with no need for radioactivity, electrophoresis, and chromatography. Specifically capturing tri-methylated H3-K79 with the detection limit as low as 0.4 ng/well and detection range from 5 ng-2 $\mu\text{g}/\text{well}$ of histone extracts. The control is conveniently included for quantification of the amount of tri-methylated H3-K79. Strip microplate format makes the assay flexible: manual or high throughput. Simple, reliable, and consistent assay conditions.