

CTLA4:B7-1[Biotinylated] Inhibitor Screening Assay Kit

Product Information

Cat.No.

Kit-1712

Product Overview

The activation of native T cells requires two signals, the specific T cell receptor recognition of MHC/Antigen on the surface of the antigen-presenting cell (APC), and the binding of B7-1 (CD80) ligand on the APC with the CD28 receptor on the T cell surface. Conversely, binding of CTLA4 to B7-1 on the T-cell surface results in an inhibitory signal and prevents T-cell activation. CTLA4:B7-1 interaction is an important drug target for the regulation of the host's response to cancer.

Size

96 reactions

Description

The CTLA4:B7-1[Biotinylated] Inhibitor Screening Assay Kit is designed for screening and profiling inhibitors of CTLA4:B7-1 signaling. The key to this kit is the high sensitivity of detection of biotin-labeled B7-1 by streptavidin-HRP. Only a few simple steps on a microtiter plate are required for the assay. First, CTLA4 is coated on a 96-well plate. Next, B7-1 is incubated with CTLA4 on the plate. Finally, the plate is treated with streptavidin-HRP followed by addition of an HRP substrate to produce chemiluminescence, which can be measured using a chemiluminescence reader.

Applications

This kit is useful for screening for inhibitors of B7-1 binding to CTLA4.

Storage

Stable at least 6 months from date of receipt, when stored as directed. Kit components require different storage conditions. Be sure to store each component at the proper temperature upon arrival.

Kit Components

B7-1, Biotin-labeled: 5 µg; -80°C CTLA4: 10 µg; -80°C Streptavidin-HRP: 15 µl; +4°C 3x CTLA4 Assay



CTLA4:B7-1[Biotinylated] Inhibitor Screening Assay Kit

Buffer: 50 ml; -20°C Blocking Buffer: 50 ml; +4°C HRP chemiluminescent substrate A (transparent bottle): 6 ml; +4°C HRP chemiluminescent substrate B (brown bottle): 6 ml; +4°C White 96-well microplate: 1; +4°C

Tel: 1-631-559-9269 1-516-512-3133

Fax: 1-631-938-8127

Email: info@creative-biomart.org

45-1 Ramsey Road, Shirley, NY 11967, USA