

Human Enolase ? Inhibitor Screening Kit (Colorimetric)

Product Information

Cat

Kit-2355

Cat.No.

Kit-2355

Product Overview

Enolase (EC 4.2.1.11), also called 2-Phospho-D-glycerate hydrolase, is a key enzyme in glycolysis. It converts 2-phosphoglycerate into phosphoenolpyruvate (PEP). It also catalyzes the reverse reaction, PEP to 2-phosphoglycerate under anabolic conditions during gluconeogenesis. In mammals, three genes encode for the Enolase α , β and γ isoforms. These isoforms combine together to form both homo and heterodimeric complexes resulting in five isozymes, three of which are commonly found in human tissues. Enolase α , also called enolase 1, exists in variety of tissues which can undergo glycolysis. Besides its role in glycolysis, increased Enolase α activity is associated with tumor invasion and metastasis; therefore, screening for novel and specific inhibitor of enolase α inhibitor may be of great interest for tumor treatment. In Human Enolase α Inhibitor Screening Kit, Enolase α catalyzes the conversion of 2-phosphoglycerate to phosphoenolpyruvate, which is subsequently used to generate an intermediate product. The intermediate product stoichiometrically reacts with OxiRed; probe generating a colorimetric signal (OD 570 nm). In the presence of Enolase α inhibitor, the reaction is impeded. An enolase α Inhibitor Control is included to compare the efficacy of the sample inhibitors. The assay is simple, fast, high-throughput adaptable and can be completed in less than 1 hr.

Applications

Screening/characterizing/studying potential inhibitors of Human Enolase α .

Storage

-20°C

Shipping

Gel Pack

Human Enolase ? Inhibitor Screening Kit (Colorimetric)

Size

100 assays

Kit Components

Enolase Assay Buffer; OxiRed Probe; Enolase Substrate; Enolase Converter; Enolase Developer; Enolase a; Enolase Inhibitor Control

Target Species

Human

Detection method Absorbance (OD 570 nm)

Features & Benefits

Rapid and easy-to-use.; High-throughput adaptable; This assay kit can be used for initial screening of compounds as anti-Enolase